665 E Vilas Road Central Point, OR 97502 Phone: (541) 779-0951 Fax: (541) 779-6498 www.crateranimalclinic.com



Robert Addington, DVM Matthew Lawrence, DVM Eric Keiser, DVM Leslie Richardson, DVM

Targeted De-worming Protocol

Due to years of rotational de-worming programs in our horse herds, many parasites have developed resistance to our de-wormers. The new recommended protocol for managing parasites in your herd encourages a fecal egg count twice yearly to identify those horses, which may be carrying, and shedding most of the parasites on your property. By targeting only those horses that need to be dewormed we can reduce the frequency in which we are de-worming, therefore decreasing the amount of parasite resistance. Please ask us if you have any questions!

March	 Fecal Egg Count on ALL HORSES in herd. Identify low, moderate and high shedders. De-worm ALL HORSES with moxidectin/praziquantel (Quest Plus) or fenbendazole (Panacur Power Pack)
June	 De-worm ALL MODERATE TO HIGH SHEDDERS with moxidectin/praziquantel (Quest Plus) or oxybendazole (Anthelcide) or Pyrantel (Strongid). No de-wormer for low shedders
September	 Optional fecal egg count on problem horses. De-worm ALL HORSES with moxidectin/praziquantel (Quest Plus) or ivermectin/praziquantel (Equimax) or fenbendazole (Panacur Power Pack)
December	 Only de-worm HIGH shedders with Oxybendazole (Anthelcide) or Pyrantel (Strongid) No de-wormer for low to moderate shedders.